## خلاصه مقالات موین گنکره عناصر کمیاب ایران، دانشگاه علوم پرشکی کاثان، ۹ و ۱۰ اسفند ۱۳۹۱

### اثرات محافظتی عنصر روی بر فعالیت غدد تیروئید و پاراتیروئید در موشهای صحرایی مسموم شده با کادمیوم

۱\* ۱ الهام مشتاقی ، پدرام ملک پوری ، احمد قورچانی

دانشجوی کارشناسی ارشد بیوشیمی، گروه بیوشیمی، دانشکده علوم پایه، دانشگاه آزاد اسلامی، واحد فلاورجان

<sup>۲</sup> دانشجوی دکترای تخصصی، گروه علوم درمانگاهی، دانشکده دامپزشکی، دانشگاه شهید چمران اهواز

<sup>۳</sup> دکترای داروسازی، گروه بیوشیمی، دانشکده داروسازی، دانشگاه علوم پزشکی اصفهان

\* نویسنده مسئول: moshtaghie.e@gmail.com

#### خلاصه:

سابقه و هدف: کادمیوم از جمله عناصر سمی است که به علت کاربردهای متعدد، از راه های گوناگون وارد بدن شده و سبب بروز عوارض مختلف می شود. هدف از انجام این پروژه بررسی اثر محافظتی روی بر کاهش تاثیرات مخرب کادمیوم بر پارامترهای سرمی مربوط به غدد تیروئید و پاراتیروئید است.

مواد و روشها: در این آزمایش تجربی ٤٠ سر موش صحرایی نر با میانگین وزنی ۳۰۳±۲۰۱ به چهارگروه پنج تایی شامل تیمار کادمیوم، روی، روی و کادمیوم، و کنترل تقسیم شدند. گروه اول روزانه سرم فیزیولوژی حاوی کادمیوم به میزان ۱ میلیگرم، گروه دوم ۰/۵ میلیگرم روی، گروه سوم ترکیبی از ۱ میلیگرم کادمیوم و ۰/۵ میلیگرم روی به ازاء هر کیلوگرم وزن بدن و گروه کنترل نیز تنها سرم فیزیولوژی دریافت نمودند. پس از پایان آزمایش در زمانهای ۳۰ و ۲۰ روز، حیوانات کشته شده و سرم خون آنها جدا شده و پارامترهای سرمی مربوط به غدد تیروئید و پاراتیروئید اندازهگیری شدند.

T3 و T3 و هورمونهای T3 و هورمونهای T3 و هورمونهای T3 و هورمونهای T3 و T3 و T3 و تتابیج به دست آمده است. نشان می دهد که کادمیوم سبب کاهش معنی دار و پاراتورمون نیز به شکل معنی داری  $(P< \cdot \cdot \cdot \circ)$  افزایش یافته است. تیمار حیوانات با روی سبب حفظ همه پارامترهای سرمی مذکور در حدود نرمال شده و یا حداقل تأثیر بازدارنده ای از خود بروز داده است.

**نتیجه گیری:** درمجموع می توان گفت در حالی که تزریق طولانی مدت کادمیوم منجر به کم کاری غده تیروئید و پرکاری غده پاراتیروئید می-شود، روی می تواند اثرات سمی این عنصر را خنثی کند.

واژ گان کلیدی: روی، کادمیوم، غده تیروئید، غده پاراتیروئید

# خلاصه مقالات سومین گنگره عناصر کمیاب ایران، دانشگاه علوم پرسنگی کاشان، ۹ و ۱۰ اسفند ۱۳۹۱

### Protective effect of zinc on serum parameters related to thyroid and parathyroid glands in rats exposed to cadmium

Moshtaghi E1\*, Malekpouri P2, Ghorchani A3

- 1- Department of Biochemistry, Faculty of Basic Sciences, Falavarjan Branch, Islamic Azad University, Isfahan, I. R. Iran.
- 2- Department of Clinical Sciences, Faculty of Veterinary Medicine, Shahid Chamran University of Ahvaz, Ahvaz, I. R. Iran.
  - 3- Department of Biochemistry, Faculty of Pharmacy, Isfahan University of Medical Science, Isfahan, I. R. Iran.

\*Corresponding author: moshtaghie.e@gmail.com

#### **Abstract:**

**Background:** Cadmium is one of the toxic metals, which can enter the body through several routes due to its numerous applications and lead to some disorders. This study aimed to investigate the protective effect of zinc on cadmium-induced damages in the serum parameters related to the thyroid and parathyroid glands.

**Materials and Methods:** In this study, 40 male rats weighting 201±30g were divided into the four separated groups (consisting of 5 rats): the cadmium treatment, zinc treatment, combination of zinc and cadmium and control groups. The normal saline containing 1 mg cadmium was injected intrapritoneally to the rats as the first treatment. The second group was treated with 0.5 mg zinc, the third one received the combined doses of cadmium (1mg) and zinc (0.5mg) and the control group received only the normal saline injection. At the end of the experiment (30 and 60 d), all rats were decapitated and sera collected to measure the serum parameters related to the thyroid and parathyroid glands.

**Results:** The obtained results showed that cadmium significantly (P<0.05) decreased the concentration of serum calcium, T3, T4 and calcitonin, but increased the Pi concentration and the activity of Paratormone and alkaline phosphatase. Zinc treatment in animals either maintained all the aforementioned serum parameters at their normal levels or at least showed a preventive-like role.

**Conclusion:** Although a long-term injection of cadmium provides hypothyroidism as well as hyperparathyroidism, zinc can be able to reduce the toxic effects of cadmium.

**Keywords:** Zinc, Cadmium, Thyroid, Parathyroid